

# The Dynamic Stall Module for FAST 8

Rick Damiani and Greg Hayman

Wednesday 13<sup>th</sup> April, 2016

## Executive Summary

The new modularization framework of FAST v.8 (Jonkman 2013) required a complete overhaul of the aerodynamics routines. AeroDyn is an aerodynamics module that can then utilize either blade element momentum theory (BEMT) or generalized dynamic wake (GDW) to calculate aerodynamic forces on blade elements. Under asymmetric conditions, such as wind shear, yawed and tilted flow, the individual blade elements undergo variations in angle of attack (AOA) that lead to unsteady aerodynamics phenomena, which can no longer be captured through the static airfoil lift and drag look-up tables. This study lays out the main theory and the organization into the modularization framework of the dynamic stall (DS) module, which includes unsteady aerodynamics under attached flow conditions and DS. DS can be called by either BEMT or GDW.

## Acknowledgments

This document was reviewed and improved thanks to the inputs by Michael Sprague, Jason Jonkman, and Bonnie Jonkman.

## Table of Contents

Acronyms . . . . .	2
List of Symbols . . . . .	2
Greek Symbols . . . . .	8
<b>1 Overview . . . . .</b>	<b>9</b>
1.1 Unsteady Attached Flow and Its Indicial Treatment . . . . .	10
1.1.1 Tangential Force . . . . .	16
1.2 TE Flow Separation . . . . .	17
1.2.1 Tangential Force . . . . .	18
1.2.2 Pitching Moment . . . . .	18
1.3 Dynamic Stall . . . . .	19
1.3.1 Tangential Force . . . . .	20
1.3.2 Pitching Moment . . . . .	21
<b>2 Inputs, Outputs, Parameters, States and Implementation of the UA . . . . .</b>	<b>22</b>
2.1 Init_Inputs . . . . .	22
2.2 Inputs u . . . . .	23
2.3 Outputs y . . . . .	23
2.4 States $x_d$ . . . . .	23
2.5 Parameters p . . . . .	24
2.6 UA Implementation . . . . .	25
2.6.1 UA_Init Routine . . . . .	25
2.6.2 UA_UpdateStates Routine . . . . .	25
2.6.2.1 DSMod Logical Flags . . . . .	25
2.6.2.2 Update Discrete States . . . . .	25
2.6.2.3 Update Other States . . . . .	26
2.6.2.3.1 $T_f$ modifications . . . . .	27
2.6.2.3.2 $T_V$ modifications . . . . .	27
2.6.2.3.3 Update 'previous time step' states . . . . .	27
2.6.3 UA_CalcOutput . . . . .	28

## List of Figures

Figure 1. Conventional stages of DS, from Leishman 2006. . . . .	10
Figure 2. Conventional stages of DS and associated $C_l$ , $C_d$ , $C_m$ as functions of AOA from Leishman 2006. . . . .	11
Figure 3. Main definitions of blade element (BE) forces (denoted via their normalized coefficients) for the unsteady aerodynamics treatment, from Damiani 2011. . . . .	12
Figure 4. Block diagram showing the order of the calls to the subroutines (1st and 2nd in the graph) and the overall organization from the parent module AeroDyn to Unsteady-Aerodynamics module (UA). . . . .	22

## List of Tables

### Acronyms

2D	two-dimensional
AOA	angle of attack
AOI	angle of incidence
BE	blade element
BEMT	blade element momentum theory
DS	dynamic stall
GDW	generalized dynamic wake
LBM	Leishman-Beddoes model
LE	leading edge
TE	trailing edge
UA	Unsteady-Aerodynamics module

### List of Symbols

$AFI_{Params}$	airfoil static tables of $C_l$ $C_d$ $C_m$ and DS parameters
$AFidx$	index pointing to the correct airfoil file/table/-database
$A_1$	constant in the expression of $\phi_\alpha^c$ and $\phi_q^c$ ; from Leishman 2011 experimental results set it to 0.3. This value is relatively insensitive for thin airfoils, but may be different for turbine airfoils. Generally speaking it should not be tuned by the user.
$A_2$	constant in the expression of $\phi_\alpha^c$ and $\phi_q^c$ ; from Leishman 2011 experimental results set it to 0.7. This value is relatively insensitive for thin airfoils, but may be different for turbine airfoils. Generally speaking it should not be tuned by the user.
$A_5$	constant in the expression of $K_q'''$ , $C_{mq}^{nc}(s, M)$ , and $k_{m,q}(M)$ ; from Leishman 2006 experimental results set it to 1
$DSMod$	switch to select handling of options and possible methods in dynamic stall

$F_R(s)$	Response Function to generic disturbance $\varepsilon(s)$
$F_R(t)$	Response Function to generic disturbance $\varepsilon(t)$
$FirstPass$	flag indicating first time step
$LESF$	leading edge separation flag
$M$	Mach Number
$NumOuts$	number of output channels
$OutFmt$	format for output data
$OutList$	list of output channels
$OutSFmt$	format for output data
$R$	Rotor Radius
$S_1$	constant in the $f$ curve best-fit for $\alpha_0 \leq \alpha \leq \alpha_1$ ; by definition it depends on the airfoil.
$S_2$	constant in the $f$ curve best-fit for $\alpha > \alpha_1$ ; by definition it depends on the airfoil.
$S_3$	constant in the $f$ curve best-fit $\alpha_2 \leq \alpha < \alpha_0$ ; by definition it depends on the airfoil.
$S_4$	constant in the $f$ curve best-fit for $\alpha < \alpha_2$ ; by definition it depends on the airfoil.
$St_{sh}$	Strouhal's shedding frequency constant, commonly taken equal to 0.19
$T'_\alpha$	Mach-dependent, non-dimensional time constant in the expression of $\phi_\alpha^{nc}$ ; it is equal to $2UT_\alpha(M)/c$
$T'_q$	Mach-dependent time constant in the expression of $\phi_q^{nc}$
$TESF$	trailing edge separation flag
$T_I$	time constant in the expression of $\phi_\alpha^{nc} = c/a_s$
$T_V$	time constant associated with the vortex lift decay process; it is used in the expression of $C_n^v$ . It depends on $Re, M$ , and airfoil class
$T_\alpha(M)$	Mach-dependent time constant in the expression of $\phi_\alpha^{nc}$
$T_f$	constant dependent on Mach, $Re$ , and airfoil shape; it is used in the expression of $D_f$ and $f''$
$T_p$	boundary-layer, leading edge (LE) pressure gradient time constant; in the expression of $D_p$ . It should be tuned based on airfoil experimental data.
$T_q(M)$	Mach-dependent time constant in the expression of $\phi_q^{nc}$
$T_{V0}$	initial value of $T_V$ .
$T_{VL}$	time constant associated with the vortex advection process; it represents the non-dimensional time in semi-chords, needed for a vortex to travel from LE to trailing edge (TE); it is used in the expression of $C_n^v$ . It depends on $Re, M$ (weakly), and airfoil. Value's range = $[6; 13]$ .
$T_{f0}$	initial value of $T_f$
$T'_{m,q}$	Mach-dependent time constant in the expression of $\phi_{m,q}^{nc}$
$T_{m,q}(M)$	Mach-dependent time constant in the expression of $\phi_{m,q}^{nc}$

$T_{sh}$	time constant associated with the vortex shedding. It allows multiple vortices to be shed at a Strouhal's frequency of 0.19.
$U$	relative air-speed
$VRTX$	vortex advection flag
$X_1$	deficiency function used in the development of $C_{n\alpha}^c(s, M)$
$X_2$	deficiency function used in the development of $C_{n\alpha}^c(s, M)$
$X_3$	deficiency function used in the development of $C_{nq}^c(s, M)$
$X_4$	deficiency function used in the development of $C_{nq}^c(s, M)$
$\Delta s$	incremental variation in $s$ for the $\Delta t$ time step
$\Delta t$	time step
$\bar{x}_{cp}$	constant in the expression of $\hat{x}_{cp}^v$ , usually = 0.2
$\alpha_{-1}$	previous time-step value of $\alpha$
$\alpha_{-2}$	two time-steps ago value of $\alpha$
$f_m''$	CENER's proposed version of lagged $f_m'$
$f_m'$	CENER's proposed lookup version of $f'$
$f''_{-1}$	previous time-step value of $f''$
$f'_{-1}$	previous time-step value of $f'$
$q_{-1}$	previous time-step value of $q$
$q_{-2}$	two time-steps ago value of $q$
$\hat{k}_1$	constant in the $C_c$ expression due to LE vortex effects
$\hat{k}_2$	constant in the $C_c$ expression due to LE vortex effects, taken equal to $2(C_n' - C_{n1}) + (f'' - f)$
$\hat{x}_{AC}$	aerodynamic center distance from LE in percent chord
$\hat{x}_{cp}^v$	center-of-pressure distance from the $1/4$ -chord, in percent chord, during the LE vortex advection process
$\hat{x}_{cp}$	center-of-pressure distance from LE in percent chord
$c$	circulatory component of the quantity at the base
$nc$	non-circulatory component of the quantity at the base
$\alpha$	relative to a step-change in $\alpha$
$n$	relative to the n-th time step
$q$	relative to a step-change in $q$
$t$	relative to the t-th time step
$a_s$	speed of sound
$b_1$	constant in the expression of $\phi_\alpha^c$ and $\phi_q^c$ ; from Leishman 2011 experimental results set it to 0.14. This value is relatively insensitive for thin airfoils, but may be different for turbine airfoils. Generally speaking it should not be tuned by the user.

$b_2$	constant in the expression of $\phi_\alpha^c$ and $\phi_q^c$ ; from Leishman 2011 experimental results set it to 0.53. This value is relatively insensitive for thin airfoils, but may be different for turbine airfoils. Generally speaking it should not be tuned by the user.
$b_5$	constant in the expression of $K_q'''$ , $C_{mq}^{nc}(s, M)$ , and $k_{m,q}(M)$ ; from Leishman 2006 experimental results set it to 5
$f'_n$	$f'$ calculated from Kirchoff's expression containing $C_c$ function of $f$
$f'_n$	$f'$ calculated from Kirchoff's expression containing $C_n$ function of $f$
$k_0$	constant in the $\hat{x}_{cp}$ curve best-fit; $= (\hat{x}_{AC} - 0.25)$
$k_1$	constant in the $\hat{x}_{cp}$ curve best-fit
$k_2$	constant in the $\hat{x}_{cp}$ curve best-fit
$k_3$	constant in the $\hat{x}_{cp}$ curve best-fit
$k_\alpha(M)$	Mach-dependent constant in the expression of $T_\alpha(M)$
$k_q(M)$	Mach-dependent constant in the expression of $T_q(M)$
$k_{m,q}(M)$	Mach-dependent constant in the expression of $T_{m,q}(M)$ and $C_{mq}^{nc}(s, M)$
$q$	non-dimensional pitching Rate $= \dot{\alpha} c / U$
$s$	non-dimensional Distance
$t$	time
$x_d$	discrete states
$D_{\alpha f, -1}$	previous time-step value of $D_{\alpha f}$
$D_{f, -1}$	previous time-step value of $D_f$
$D_{p, -1}$	previous time-step value of $D_p$
$K'_{\alpha, -1}$	previous time-step value of $K'_\alpha$
$K'''_{q, -1}$	previous time-step value of $K'''_q$
$K''_{q, -1}$	previous time-step value of $K''_q$
$K'_{q, -1}$	previous time-step value of $K'_q$
$X_{1, -1}$	previous time-step value of $X_1$
$X_{2, -1}$	previous time-step value of $X_2$
$X_{3, -1}$	previous time-step value of $X_3$
$X_{4, -1}$	previous time-step value of $X_4$
$C_n^{pot, -1}$	previous time-step value of $C_n^{pot}$
$C_n^v, -1$	previous time-step value of $C_n^v$
$C_V, -1$	previous time-step value of $C_V$
$c$	chord length
$C_c$	2D tangential (along chord) force coefficient
$C_c^{fs}$	2D tangential (along chord) force coefficient under separated TE flow separation conditions.
$C_c^{pot}$	2D along-chord force coefficient under attached (potential) flow conditions
$C_d$	2D drag coefficient
$C_{d0}$	2D drag coefficient at 0-lift



$C_l$	2D lift coefficient
$C_{l\alpha}$	slope of the 2D lift coefficient curve
$C_m$	2D pitching moment coefficient about 1/4-chord, positive if nose up
$C_{m0}$	2D pitching moment coefficient at 0-lift, positive if nose up
$C_{m\alpha}(s, M)$	pitching moment coefficient response to step change in $\alpha$
$C_{m\alpha}^c(s, M)$	circulatory component of the pitching moment coefficient response to step change in $\alpha$
$C_{m\alpha}^{nc}(s, M)$	non-circulatory component of the pitching moment coefficient response to step change in $\alpha$
$C_{m\alpha,q}(s, M)$	moment coefficient response to step change in $\alpha$ and $q$
$C_{m\alpha,q}^c(s, M)$	circulatory component of $C_{m\alpha,q}(s, M)$
$C_{m\alpha,q}^{nc}(s, M)$	non-circulatory component of $C_{m\alpha,q}(s, M)$
$C_{mq}(s, M)$	pitching moment coefficient response to step change in $q$
$C_{mq}^c(s, M)$	circulatory component of the pitching moment coefficient response to step change in $q$
$C_{mq}^{nc}(s, M)$	non-circulatory component of the moment coefficient response to step change in $q$
$C_{m\alpha}$	slope of the 2D pitching moment coefficient curve
$C_m^{fs}$	2D tangential 1/4-chord pitching moment coefficient under separated TE flow separation conditions.
$C_m^{pot}$	2D moment coefficient under attached (potential) flow conditions about 1/4-chord location
$C_{mq}(s, M)$	slope of the pitching moment coefficient vs. $q$ curve
$C_m^v$	pitching moment coefficient due to the presence of LE vortex
$C_n$	2D normal-to-chord force coefficient
$C_{n1}$	critical value of $C_n'$ at LE separation. It should be extracted from airfoil data at a given Mach and Reynolds number. It can be calculated from the static value of $C_n$ at either the break in the pitching moment or the loss of chord force at the onset of stall. It is close to the condition of maximum lift of the airfoil at low Mach numbers.
$C_{n2}$	critical value of $C_n'$ at LE separation for negative AOAs; analogous to $C_{n1}$ .
$C_{n\alpha}(s, M)$	normal force coefficient response to step change in $\alpha$
$C_{n\alpha}^c(s, M)$	circulatory component of the normal force coefficient response to step change in $\alpha$
$C_{n\alpha}^{nc}(s, M)$	non-circulatory component of the normal force coefficient response to step change in $\alpha$
$C_{n\alpha,q}(s, M)$	normal force coefficient response to step change in $\alpha$ and $q$
$C_{n\alpha,q}^c(s, M)$	circulatory component of $C_{n\alpha,q}(s, M)$

$C_{n\alpha,q}^{nc}(s,M)$	non-circulatory component of $C_{n\alpha,q}(s,M)$
$C_n^c(s,M)$	circulatory component of $C_{n\alpha,q}(s,M)$
$C_{nq}(s,M)$	normal force coefficient response to step change in $q$
$C_{nq}^c(s,M)$	circulatory component of the normal force coefficient response to step change in $q$
$C_{nq}^{nc}(s,M)$	non-circulatory component of the normal force coefficient response to step change in $q$
$C_{n\alpha}$	slope of the 2D normal coefficient curve, should be similar to $C_{l\alpha}$
$C_{n\alpha}^c(s,M)$	slope of the circulatory normal force coefficient vs. $\alpha$ curve
$C_n^{fs}$	normal force coefficient under separated TE flow separation conditions.
$C_n'$	lagged component of $C_n$ in the TE separated treatment
$C_n^{pot}$	2D normal-to-chord force coefficient under attached (potential) flow conditions
$C_n^{pot,c}$	circulatory part of 2D normal-to-chord force coefficient under attached (potential) flow conditions
$C_{nq}(s,M)$	slope of the normal force coefficient vs. $q$ curve
$C_n^v$	normal force coefficient due to the presence of LE vortex
$C_V$	contribution to the normal force coefficient due to accumulated vorticity in the LE vortex
$D_{\alpha f}$	deficiency function for $\alpha_f$
$D_f$	deficiency function for $f'$
$D_p$	deficiency function for $C_n'$
$f$	separation point distance from LE in percent chord
$f_m$	CENER's proposed version of $f$ extracted from the $C_m$ static tables, assuming $C_m = C_n f_m$
$f_m'$	version of $f_m$ extracted from the airfoil $C_m$ static tables with $\alpha_f$ as input parameter
$f_m''$	lagged version of $f_m'$
$f'$	separation point distance from LE in percent chord under unsteady conditions
$f''$	lagged version of $f'$ accounting for unsteady boundary layer response
$k$	reduced frequency
$K_\alpha$	backward finite difference of $\alpha$ at the t-th time step
$K_\alpha'$	deficiency function for $C_{n\alpha}^{nc}(s,M)$
$K_q$	backward finite difference of $q$ at the t-th time step
$K_q'$	deficiency function for $C_{nq}^{nc}(s,M)$
$K_q''$	deficiency function for $C_{m_q}^{nc}(s,M)$
$K_q'''$	deficiency function for $C_{m_q}^c(s,M)$
$Re$	airfoil-chord Reynolds Number

$U$  air velocity magnitude relative to the airfoil

## Greek Symbols

$\Delta_{\alpha 0}$	$\alpha - \alpha_0$
$\alpha_0$	0-lift angle of attack
$\alpha_1$	angle of attack at $f=0.7$ , (approximately the stall angle) for $\alpha \geq \alpha_0$
$\alpha_2$	angle of attack at $f=0.7$ , for $\alpha < \alpha_0$
$\alpha_e$	effective angle of attack at $3/4$ -chord
$\alpha$	angle of attack
$\varepsilon(s)$	generic disturbance
$\varepsilon(t)$	generic disturbance
$\eta_e$	recovery factor $\simeq [0.85 - 0.95]$ to account for viscous effects at limited or no separation on $C_c$
$\omega$	generic frequency
$\phi(s, M)$	indicial response function
$\phi(t, M)$	indicial response function
$\phi_\alpha^c$	normal force coefficient, circulatory indicial response function to a step change in $\alpha$
$\phi_\alpha^{nc}$	normal force coefficient, non-circulatory indicial response function to a step change in $\alpha$
$\phi_q^c$	normal force coefficient, circulatory indicial response function to a step change in $q$
$\phi_q^{nc}$	normal force coefficient, non-circulatory indicial response function to a step change in $q$
$\phi_{m,\alpha}^{nc}$	pitching moment coefficient, non-circulatory indicial response function to a step change in $\alpha$
$\phi_{m,q}^c$	pitching moment coefficient, circulatory indicial response function to a step change in $q$
$\phi_{m,q}^{nc}$	pitching moment coefficient, non-circulatory indicial response function to a step change in $q$
$\sigma_1$	generic multiplier for $T_f$
$\sigma_3$	generic multiplier for $T_V$
$\sigma_s$	generic integrand coordinate
$\sigma_l$	generic integrand coordinate
$\tau_V$	time variable, tracking the travel of the LE vortex over the airfoil suction surface. It is made dimensionless via the semi-chord: $\tau_V = t * 2U / c$ . If less than $2T_{VL}$ , it renders $VRTX=True$ ; if less than $T_{VL}$ then the vortex is still on the airfoil.
$\alpha_f$	effective angle of incidence (AOI) which would give the same unsteady LE pressure gradient under static conditions; used to calculate $f'$
$\alpha'_f$	lagged version of $\alpha_f$ ; used in Minnema 1998 calculation of $C_m$ under separated conditions
$\beta_M$	Prandtl-Glauert compressibility correction factor $\sqrt{1 - M^2}$

# 1 Overview

The main theory follows the work by Leishman and Beddoes 1986, 1989; Pierce and Hansen 1995; Pierce 1996; Leishman 2011; Damiani 2011. Dynamic stall is a well-known phenomenon that can affect wind turbine performance and loading especially during yawed operations, and that can result in large unsteady stresses on the structures.

Dynamic stall manifests as a delay in the onset of flow separation to higher AOAs that would otherwise occur under static (steady) conditions, followed by an abrupt flow separation from the LE of the airfoil Leishman 2011. The LE separation is the fundamental characteristic of the DS of an airfoil; in contrast, quasi-steady stall would start from the airfoil TE.

DS occurs for reduced frequencies above 0.02.

$$k = \frac{\omega c}{2U} \quad (1.1)$$

The five stages of DS are as follows and shown in Fig. 1 and Fig. 2:

1. Onset of flow reversal
2. Flow separation and vorticity accumulation at the LE
3. Shedding of the vortex and convection along the suction surface of the airfoil (lift increases)
4. Lift Stall: vortex is shed in the wake and lift abrupt drop-off
5. Re-attachment of the flow at AOAs considerably lower than static AOAs (hysteresis)

The model chosen to represent unsteady aerodynamics and dynamic stall is the Leishman-Beddoes model (LBM), because it is the most widely used and has the most available support throughout the community and it has shown reasonable success when compared to experimental data. The LBM is a postdictive model, and as such it will not solve equations of motion, though the principles are fully rooted in the physics of unsteady flow.

In the LBM, the different processes are modeled as first-order subsystems with differential equations with pre-determined constants to match experimental results. Therefore, knowledge of the airfoil characteristics under unsteady aerodynamics is a prerogative of the LBM. The LBM may also be described as an indicial response (i.e., response to a series of small disturbances) model for attached flow, extended to account for separated flow effects and vortex lift. Forces are computed as normal and tangential (to chord) and pitching moment about the 1/4-chord location, see also Fig. 3 and Eq. (1.2).


$$\begin{aligned} C_l &= C_n \cos \alpha + C_c \sin \alpha \\ C_d &= C_n \sin \alpha - C_c \cos \alpha + C_{d0} \end{aligned} \quad (1.2)$$

$$\begin{aligned} C_n &= C_l \cos \alpha + (C_d - C_{d0}) \sin \alpha \\ C_c &= C_l \sin \alpha - (C_d - C_{d0}) \cos \alpha \end{aligned} \quad (1.3)$$

The original model was developed for helicopters, but it has been successfully applied to wind turbines (see Pierce 1996; Gupta and Leishman 2006). Yawed flow conditions, Coriolis and centrifugal forces that lead to three-dimensional effects were not included in the original model.

Unsteady aerodynamics is mostly driven by 2D flow aspects, where the time scale is on the order of tenths of seconds, or  $\sim c/\omega R$ .

The LBM considers a number of unsteady aerodynamics conditions, namely: attached flow conditions and TE separation before stall; delays and lags associated with the unsteady onset of dynamic stall and accompanying boundary



**Figure 1. Conventional stages of DS, from Leishman 2006.**

layer development; advection of the LE vortex, shedding in the wake, and suppression of TE separation in favor of LE separation. The LBM can be subdivided into three main submodules:

1. Unsteady, Attached Flow Solution via Indicial Treatment (potential flow)
2. TE Flow Separation
3. DS and Vorticity Advection

## 1.1 Unsteady Attached Flow and Its Indicial Treatment

The advantage of the indicial treatment is that a response to an arbitrary forcing can be obtained through superposition of response-functions to a step variation in AOA, in pitch rate, or in heave (plunging) motion. The superposition is carried out via the so-called Duhamel Integral Leishman 2006, which for the generic response  $F_R(t)$  to a generic disturbance  $\varepsilon(t)$  can be written as in Eq. (1.4):

$$\begin{aligned}
 F_R(t) &= \varepsilon(0)\phi(t, M) + \int_0^t \frac{d\varepsilon}{d\sigma_t}(\sigma_t)\phi(t - \sigma_t, M)d\sigma_t \\
 \text{or} \\
 F_R(s) &= \varepsilon(0)\phi(s, M) + \int_0^s \frac{d\varepsilon}{d\sigma_s}(\sigma_s)\phi(s - \sigma_s, M)d\sigma_s
 \end{aligned} \tag{1.4}$$

where the non-dimensional distance,  $s$ , is defined as:

$$s = \frac{2}{c} \int_0^t U(t)dt \tag{1.5a}$$

$$\Delta s = \frac{2}{c} U(t)\Delta t \tag{1.5b}$$

where the airfoil half chord is considered as the non-dimensionalizing factor ( $c/2$ ).

The indicial functions are surmised into two components: the first is related to the non-circulatory loading (piston theory and acoustic wave theory); the second originates from the development of circulation about the airfoil.





Figure 2. Conventional stages of DS and associated  $C_l$ ,  $C_d$ ,  $C_m$  as functions of AOA from Leishman 2006.



**Figure 3. Main definitions of BE forces (denoted via their normalized coefficients) for the unsteady aerodynamics treatment, from Damiani 2011.**

The non-circulatory part depends on the instantaneous airfoil motion, but also on the time history of the prior motion. The circulatory response can be calculated via the 'lumped approach', where the effects of step changes in AOA, pitch rate, heave motion, etc., are combined into an effective AOA at the  $3/4$ -chord station.

The normal force coefficient response to a step-change in non-dimensional pitch rate  $q$  and a step-change in AOA can be written as a function of the indicial functions as shown in Eq. (1.6):

$$\begin{aligned} C_{n_{\alpha,q}}(s, M) &= C_{n_{\alpha}}(s, M) + C_{n_q}(s, M) = C_{n_{\alpha}}\alpha + C_{n_q}(s, M)q \\ C_{n_{\alpha}}(s, M) &= \frac{4}{M}\phi_{\alpha}^{nc} + \frac{C_{n_{\alpha}}}{\beta_M}\phi_{\alpha}^c \\ C_{n_q}(s, M) &= \frac{1}{M}\phi_q^{nc} + \frac{C_{n_{\alpha}}}{2\beta_M}\phi_q^c \end{aligned} \quad (1.6)$$

Analogously the pitching moment coefficient about the  $1/4$ -chord can be derived via indicial response as shown in Eq.(1.7):

$$\begin{aligned} C_{m_{\alpha,q}}(s, M) &= C_{m_{\alpha}}(s, M) + C_{m_q}(s, M) = C_{m_{\alpha}}\alpha + C_{m_q}(s, M)q \\ C_{m_{\alpha}}(s, M) &= -\frac{1}{M}\phi_{m,\alpha}^{nc} - \frac{C_{m_{\alpha}}}{\beta_M}\phi_{\alpha}^c (\hat{x}_{AC} - 0.25) + C_{m0} \\ C_{m_q}(s, M) &= -\frac{7}{12M}\phi_{m,q}^{nc} - \frac{C_{m_{\alpha}}}{16\beta_M}\phi_{m,q}^c \end{aligned} \quad (1.7)$$

where  $C_{m0}$  is positive if causes a pitch up of the airfoil, as seen in Fig. 3. Also note that the circulatory component of the pitching moment response to a step-change in  $\alpha$  is a function of the  $C_{n_{\alpha}}^c(s, M)$ .

The non-dimensional pitch-rate  $q$  is given by:

$$\begin{aligned} q &= \frac{\dot{\alpha}c}{U} \simeq K_{\alpha t}c/U \\ \text{with: } K_{\alpha t} &= \frac{\alpha_t - \alpha_{t-1}}{\Delta t} \end{aligned} \quad (1.8)$$

For small  $\Delta t$ s, the finite difference  $K_{\alpha}$  can be subjected to significant numerical noise. In order to smooth the terms associated with derivatives, a low-pass filter is introduced. The filter is applied to  $\alpha$ ,  $q$ , and its derivative  $K_q$  (which is

used later) as defined in Eq. (1.9).

$$\begin{aligned}
\alpha_{LP_t} &= C_{LP}\alpha_{t-1} + (1 - C_{LP})\alpha_t \\
q &= \frac{(\alpha_{LP_t} - \alpha_{LP_{t-1}})c}{U\Delta t} \\
q_{LP} &= C_{LP}q_{LP_{t-1}} + (1 - C_{LP})q_t \\
K_{\alpha LP} &= \frac{q_{LP}U}{c} \\
K_{q_t} &= \frac{q_t - q_{t-1}}{\Delta t} \\
K_{q_{LP_t}} &= C_{LP}K_{q_{LP_{t-1}}} + (1 - C_{LP})K_{q_t}
\end{aligned} \tag{1.9}$$

where :

$$C_{LP} = e^{-2\pi\Delta t\zeta_{LP}}$$

where  $\alpha_{LP_t}$  is the low-pass filtered value of  $\alpha$ ;  $q_{LP}$  is the low-pass filtered value of  $q$ ;  $K_{\alpha LP}$  is the modified value value of  $K_{\alpha}$  due to filtered  $\alpha$  and  $q$ ;  $K_{q_t}$  is the backward finite difference of  $q$  at the  $t$ -th time step;  $K_{q_{LP_t}}$  is the low-pass filtered value of  $K_{q_t}$ ;  $C_{LP}$  is the low-pass filter constant; and  $\zeta_{LP}$  is the low-pass filter frequency cutoff  $-3$  dB).

From here on, the 'LP' subscript is dropped, and with the understanding that quantities such as  $\alpha$ ,  $K_{\alpha}$ ,  $q$ , and  $K_q$  denote the respective filtered quantities  $\alpha_{LP_t}$ ,  $K_{\alpha LP}$ ,  $q_{LP}$ ,  $K_{q_{LP_t}}$  as defined in Eq. (1.9).

The indicial responses can then be approximated as in Eq. (1.10) (Leishman and Beddoes 1989; Johansen 1999):

$$\begin{aligned}
\phi_{\alpha}^c &= \phi_q^c = 1 - A_1 \exp(-b_1 \beta_M^2 s) - A_2 \exp(-b_2 \beta_M^2 s) \\
\phi_{\alpha}^{nc} &= \exp\left(-\frac{s}{T'_{\alpha}}\right) \\
\phi_q^{nc} &= \exp\left(-\frac{s}{T'_q}\right) \\
\phi_{m,q}^{nc} &= \exp\left(-\frac{s}{T'_{m,q}}\right)
\end{aligned} \tag{1.10}$$

One could find analogous expressions for  $\phi_{m,q}^c, \phi_{m,\alpha}^{nc}$ , but they are not shown here because further simplified expressions will be derived below.

By making use of exact results for short times  $0 \leq s \leq 2M/(M+1)$  (Lomax et al. 1952), Leishman 2011 shows that:

$$\begin{aligned}
T_{\alpha}(M) &= 0.75 \frac{c}{2U} T'_{\alpha} = 0.75 \frac{c}{2Ma_s} T'_{\alpha} = 0.75 k_{\alpha}(M) T_I \\
T_q(M) &= 0.75 \frac{c}{2U} T'_q = 0.75 \frac{c}{2Ma_s} T'_q = 0.75 k_q(M) T_I
\end{aligned} \tag{1.11}$$

where:

$$k_{\alpha}(M) = \left[ (1-M) + \frac{C_{n\alpha}}{2} M^2 \beta_M (A_1 b_1 + A_2 b_2) \right]^{-1} = \left[ (1-M) + \frac{C_{n\alpha}}{2} M^2 \beta_M 0.413 \right]^{-1} \tag{1.12a}$$

$$k_q(M) = \left[ (1-M) + C_{n\alpha} M^2 \beta_M (A_1 b_1 + A_2 b_2) \right]^{-1} = \left[ (1-M) + C_{n\alpha} M^2 \beta_M 0.413 \right]^{-1} \tag{1.12b}$$

$$T_I = \frac{c}{a_s} \tag{1.12c}$$



Note that Leishman 2011 recommends the use of the factor 0.75 for  $T_\alpha(M)$  and  $T_q(M)$  to account for three-dimensional effects not included in piston theory.

The values of the  $A_1$ - $b_2$  constants are independent of  $M$  and are determined from experimental data on oscillating airfoils in wind tunnels.

For the circulatory component of the aerodynamic force response, the lumped approach can lead to a direct solution of  $C_{n\alpha,q}^c(s, M)$ . Considering the circulatory part  $C_{n\alpha}^c(s, M)$  of Eq. (1.6) for the response to the step in  $\alpha$ , one can write:

$$C_{n\alpha}^c(s, M) = \int_{s_0}^s \frac{C_{n\alpha}}{\beta_M} \phi_\alpha^c \alpha(s) ds \simeq C_{n\alpha}^c(s, M) \Delta\alpha \quad (1.13)$$

where  $C_{n\alpha}^c(s, M) = \frac{C_{n\alpha}}{\beta_M}$

By using Eq. (1.4) with  $\phi(s, M)$  replaced by  $\phi_\alpha^c$  and  $\varepsilon(s)$  by  $\alpha$ , Eq. (1.13) rewrites:

$$C_{n\alpha,q}^c(s, M) = C_{n\alpha}^c(s, M) \left[ \alpha(s_0) \phi_\alpha^c(s) + \int_{s_0}^s \frac{d\alpha}{d\sigma_s} (\sigma_s) \phi_\alpha^c(s - \sigma_s, M) d\sigma_s \right] = C_{n\alpha}^c(s, M) \alpha_e \quad (1.14)$$

where  $\alpha_e$  is an effective angle of attack at  $3/4$ -chord accounting for a step variation in  $\alpha$ , pitching rate, heave, and velocity (lumped approach). By applying the first of Eq. (1.10), and setting  $s_0 = 0$ , Eq. (1.14) can be simplified to arrive at an expression for  $\alpha_e$  at the  $n$ -th time step, i.e.,  $\alpha_{e_n}$ :

$$\alpha_{e_n}(s, M) = (\alpha_n - \alpha_0) - X_{1n}(\Delta s) - X_{2n}(\Delta s) \quad (1.15)$$

where the  $\int_{s_0}^s [\dots] d\sigma_s$  was divided into two steps considering a distance interval  $\Delta s$ , i.e.,  $\int_0^s [\dots] d\sigma_s$  and  $\int_s^{s+\Delta s} [\dots] d\sigma_s$ , by carrying out the algebra a recursive expression for  $X_1$  and  $X_2$  can be found:

$$\begin{aligned} X_{1n} &= X_{1n-1} \exp(-b_1 \beta_M^2 \Delta s) + A_1 \exp(-b_1 \beta_M^2 \frac{\Delta s}{2}) \Delta \alpha_n \\ X_{2n} &= X_{2n-1} \exp(-b_2 \beta_M^2 \Delta s) + A_2 \exp(-b_2 \beta_M^2 \frac{\Delta s}{2}) \Delta \alpha_n \end{aligned} \quad (1.16)$$

Note that  $\alpha_0$  was introduced into Eq.(1.15), because  $\alpha_e$  is an effective AOI and not AOA.

Similarly to the above development, the circulatory contribution to  $C_{n\alpha,q}^c(s, M)$  from a step change in  $q$  can be derived as:

$$C_{nq}^c(s, M) = \frac{C_{n\alpha}^c(s, M)}{2} (q_n) - X_{3n}(\Delta s) - X_{4n}(\Delta s) \quad (1.17)$$

with

$$\begin{aligned} X_{3n} &= X_{3n-1} \exp(-b_1 \beta_M^2 \Delta s) + A_1 \exp(-b_1 \beta_M^2 \frac{\Delta s}{2}) \Delta q \\ X_{4n} &= X_{4n-1} \exp(-b_2 \beta_M^2 \Delta s) + A_2 \exp(-b_2 \beta_M^2 \frac{\Delta s}{2}) \Delta q \end{aligned} \quad (1.17a)$$

Eqs. (1.17)-(1.17a) are used by González 2014.

However, following the original LBM method, the lumped approach can account for any effect to  $\alpha$ , including step changes in  $q$ , so Eq. (1.17) is not necessary, and it is virtually included via Eq. (1.14) and (1.15).

The non-circulatory part cannot be handled via the superposition (lumped approach), therefore the contribution from step changes in  $\alpha$  and  $q$  need to be kept separate:

$$C_{n_{\alpha,q}}^{nc}(s, M) = C_{n_{\alpha}}^{nc}(s, M) + C_{n_q}^{nc}(s, M) \quad (1.18)$$

Now, using Duhamel's integral (1.4) on the non-circulatory component  $C_{n_{\alpha}}^{nc}(s, M)$  (see Eq.(1.18)) with the  $\phi_{\alpha}^{nc}$  from Eq.(1.10), one can arrive at:

$$\begin{aligned} C_{n_{\alpha}}^{nc}(s, M) &= \frac{4T_{\alpha}(M)}{M} (K_{\alpha t} - K'_{\alpha t}) \\ K'_{\alpha t} &= K'_{\alpha t-1} \exp\left(-\frac{\Delta t}{T_{\alpha}(M)}\right) + (K_{\alpha t} - K_{\alpha t-1}) \exp\left(-\frac{\Delta t}{2T_{\alpha}(M)}\right) \end{aligned} \quad (1.19)$$

Note that in Eq. (1.19),  $K'_{\alpha}$  is the deficiency function for  $C_{n_{\alpha}}^{nc}(s, M)$ .

For  $C_{n_q}^{nc}(s, M)$ , an analogous procedure leads to:

$$\begin{aligned} C_{n_q}^{nc}(s, M) &= -\frac{T_q(M)}{M} (K_{q t} - K'_{q t}) \\ K'_{q t} &= K'_{q t-1} \exp\left(-\frac{\Delta t}{T_q(M)}\right) + (K_{q t} - K_{q t-1}) \exp\left(-\frac{\Delta t}{2T_q(M)}\right) \end{aligned} \quad (1.20)$$

So finally, the expression for the total normal force under attached conditions can be expressed as:

$$\begin{aligned} C_n^{pot} &= C_n^{pot,c} + C_n^{pot,nc} \\ C_n^{pot} &= C_{n_{\alpha,q}}^c(s, M) + C_{n_{\alpha,q}}^{nc}(s, M) = C_{n_{\alpha}}^c(s, M)\alpha_e + \frac{4T_{\alpha}(M)}{M} (K_{\alpha t} - K'_{\alpha t}) + \frac{T_q(M)}{M} (K_{q t} - K'_{q t}) \end{aligned} \quad (1.21)$$

with

$$\begin{aligned} C_n^{pot,c} &= C_{n_{\alpha,q}}^c(s, M) = C_{n_{\alpha}}^c(s, M)\alpha_e \\ C_n^{pot,nc} &= C_{n_{\alpha,q}}^{nc}(s, M) = \frac{4T_{\alpha}(M)}{M} (K_{\alpha t} - K'_{\alpha t}) + \frac{T_q(M)}{M} (K_{q t} - K'_{q t}) \end{aligned}$$

Now turning back to the pitching moment, following Johansen 1999 the circulatory component  $C_{m_q}^c(s, M)$  can be written as:

$$C_{m_q}^c(s, M) = -\frac{C_{n_{\alpha}}}{16\beta_M} (q - K_q''') \frac{c}{U} \quad (1.22)$$

where

$$K_q''' = K_q'''_{t-1} \exp(-b_5 \beta_M^2 \Delta s) + A_5 \Delta q_t \exp\left(-b_5 \beta_M^2 \frac{\Delta s}{2}\right) \quad (1.23)$$

The non-circulatory component of the pitching moment response to step-change in  $\alpha$ ,  $C_{m_{\alpha}}^{nc}(s, M)$ , writes (Leishman and Beddoes 1986; Johansen 1999):

$$C_{m_{\alpha}}^{nc}(s, M) = -\frac{1}{M} \phi_{m,\alpha}^{nc} = -\frac{C_{n_{\alpha}}^{nc}(s, M)}{4} \quad (1.24)$$

which implies

$$\phi_{m,\alpha}^{nc} = \phi_{\alpha}^{nc} \quad (1.25)$$

The other component  $C_{m_q}^{nc}(s, M)$  writes (Leishman 2006):

$$\begin{aligned} C_{m_q}^{nc}(s, M) &= -\frac{7}{12M} \phi_{m,q}^{nc} = -\frac{7k_{m,q}(M)^2 T_I}{12M} (K_q - K_q'') \\ \text{with:} & \\ k_{m,q}(M) &= \frac{7}{15(1-M) + 1.5C_{n\alpha} A_5 b_5 \beta_M M^2} \\ K_{q_t}'' &= K_{q_{t-1}}'' \exp\left(-\frac{\Delta t}{k_{m,q}(M)^2 T_I}\right) + (K_{q_t} - K_{q_{t-1}}) \exp\left(-\frac{\Delta t}{2k_{m,q}(M)^2 T_I}\right) \end{aligned} \quad (1.26)$$

where the same procedure as before was used to arrive at a deficiency function using Duhamel's integral (Eq. (1.4)) and equating the expressions of the  $C_{m_{\alpha,q}}(s, M)$  at  $s \rightarrow 0$  (see Leishman 2006).

Finally, the expression for the total pitching moment at  $1/4$ -chord under attached conditions can be expressed as:

$$\begin{aligned} C_m^{pot} &= C_{m_{\alpha,q}}^c(s, M) + C_{m_{\alpha,q}}^{nc}(s, M) = \\ &C_{m0} - \frac{C_{n\alpha}}{\beta_M} \phi_{\alpha}^c (\hat{x}_{AC} - 0.25) + \\ &-\frac{C_{n\alpha}}{16\beta_M} (q - K_q''') \frac{c}{U} + \\ &-\frac{T_{\alpha}(M)}{M} (K_{\alpha_t} - K_{\alpha_t}') + \\ &-\frac{7k_{m,q}(M)^2 T_I}{12M} (K_q - K_q'') \end{aligned} \quad (1.27)$$

Note that the pitching moment treatment is slightly different from what is in AeroDyn v13 and Damiani 2011, and it is more in line with Leishman 2006 and Johansen 1999. If Minnema 1998 suggestions is used, then the  $C_{m_q}^{nc}(s, M)$  (last term in Eq. (1.27)) is to be replaced by:

$$C_{m_q}^{nc}(s, M) = -\frac{7}{12M} \phi_{m,q}^{nc} = -\frac{C_{n_q}^{nc}(s, M)}{4} - \frac{k_{\alpha}(M)^2 T_I}{3M} (K_q - K_q'') \quad (1.28)$$

### 1.1.1 Tangential Force

The tangential force along the chord can be written as in Eq. (1.29) from Leishman 2011:

$$C_c^{pot} = C_n^{pot,c} \tan(\alpha_e + \alpha_0) \quad (1.29)$$

In potential flow, D'Alembert's paradox leads to the absence of drag; therefore, from Eq. (1.3),  $C_l \cos \alpha = C_n$ , and  $C_c = C_l \sin \alpha$  which bring forth Eq.(1.29). Since  $\alpha_e$  is a virtual AOI at  $3/4$ -chord, we needed to add the  $\alpha_0$ . Since this drag treatment has roots only in the circulatory lift derivation, the non-circulatory part is dropped as seen in Eq. (1.29).

## 1.2 TE Flow Separation

The base of this dynamic system is Kirchoff's theory, which can be expressed as follows (Leishman 2006):

$$\begin{aligned} C_n(\alpha, f, s, M) &= C_{n\alpha}^c(s, M) (\alpha - \alpha_0) \left( \frac{1+\sqrt{f}}{2} \right)^2 \\ C_c(\alpha, f, s, M) &= \eta_e C_{n\alpha}^c(s, M) (\alpha - \alpha_0) \sqrt{f} \tan(\alpha) \end{aligned} \quad (1.30)$$

with  $f$  the separation point distance from LE in percent chord, and  $\eta_e$  the recovery factor  $\simeq [0.85 - 0.95]$  to account for viscous effects at limited or no separation on  $C_c$ .

If the airfoil's  $C_l, C_d$ , and  $C_m$  characteristics are known, then Eq.1.30 may be solved for  $f$ . Leishman 2011 suggests the use of best-fit curves obtained from static measurements on airfoils, of the type:

$$f = \begin{cases} 1 - 0.3 \exp\left(\frac{\alpha - \alpha_1}{S_1}\right), & \text{if } \alpha_0 \leq \alpha \leq \alpha_1 \\ 1 - 0.3 \exp\left(\frac{\alpha_2 - \alpha}{S_3}\right), & \text{if } \alpha_2 \leq \alpha < \alpha_0 \\ 0.04 + 0.66 \exp\left(\frac{\alpha_1 - \alpha}{S_2}\right), & \text{if } \alpha > \alpha_1 \\ 0.04 + 0.66 \exp\left(\frac{\alpha - \alpha_2}{S_4}\right), & \text{if } \alpha < \alpha_2 \end{cases} \quad (1.31)$$

$S_1$ - $S_2$  (and the analogous  $S_3$ - $S_4$  for  $\alpha < \alpha_0$ ) are best-fit constants that define the abruptness of the static stall.  $\alpha_1$  is the angle of attack at  $f=0.7$ , (approximately the stall angle) for  $\alpha \geq \alpha_0$ , whereas  $\alpha_2$  is the angle of attack at  $f=0.7$ , for  $\alpha < \alpha_0$ .

Now accounting for unsteady conditions, the TE separation point gets modified due to temporal effects on airfoil pressure distribution and boundary layer response. LE separation occurs when a critical pressure at the LE, corresponding to a critical value of the normal force  $C_{n1}$ , is reached. The circulatory normal force needs to be modified to account for the lagged boundary layer response. In order to arrive at a new expression for  $C_n$ , we start by accounting for the separation point location under unsteady conditions, which can be calculated starting from an effective AOI,  $\alpha_f$ :

$$\alpha_f = \frac{C_n'}{C_{n\alpha}^c(s, M)} + \alpha_0 \quad (1.32)$$

where an effective  $C_n'$  is used, calculated as in Eq. (1.33):

$$\begin{aligned} C_n' &= C_n^{pot} - D_p \\ D_{p_t} &= D_{p_{t-1}} \exp\left(-\frac{\Delta s}{T_p}\right) + (C_n^{pot} - C_n^{pot_{t-1}}) \exp\left(-\frac{\Delta s}{2T_p}\right) \end{aligned} \quad (1.33)$$

Note  $T_p$  is boundary-layer, LE pressure gradient time constant; in the expression of  $D_p$ . It should be tuned based on airfoil experimental data. and is an empirically based quantity that should be tuned from experimental data. Johansen 1999 employs two time constants  $T_{p\alpha}$  and  $T_{pq}$  as the  $C_n'$  is separated into two contributions, one from  $\alpha$  and from  $q$ .

Given the new  $C_n'$ , one can attain a new formulation for  $f$ , i.e.,  $f''$ , which accounts for delays in the boundary layer, and that will be used via Kirchoff's treatment to arrive at the new  $C_n$ :

$$\begin{aligned} f'' &= f' - D_f \\ D_{f_t} &= D_{f_{t-1}} \exp\left(-\frac{\Delta s}{T_f}\right) + (f'_t - f'_{t-1}) \exp\left(-\frac{\Delta s}{2T_f}\right) \end{aligned} \quad (1.34)$$

where  $f'$  is the separation point distance from LE in percent chord under unsteady conditions that can be obtained from the best-fit in Eq. (1.31) replacing  $\alpha$  with  $\alpha_f$ .

**Alternatively**,  $f'$  can be derived from a direct lookup table of static airfoil data reversing Eq. (1.30). In fact, two values of  $f'$  could be calculated: one for  $C_n$  and one for  $C_c$ .

Also note that  $T_f$  is a Mach,  $Re$  and airfoil dependent time constant associated with the motion of the separation point along the suction surface of the airfoil. constant dependent on Mach,  $Re$ , and airfoil shape; it is used in the expression of  $D_f$  and  $f''$  gets modified via multipliers ( $\sigma_1$ ) depending on the phase of the separation or reattachment as discussed later; here it suffices noting that  $T_f$  can be written as a modified version of the initial value  $T_{f0}$ :

$$T_f = T_{f0}/\sigma_1 \quad (1.35)$$

Finally, the normal force coefficient  $C_n^{fs}$ , after accounting for separated flow from the TE becomes:

$$C_n^{fs} = C_{n\alpha,q}^{nc}(s, M) + C_{n\alpha,q}^c(s, M) \left( \frac{1 + \sqrt{f''}}{2} \right)^2 = C_{n\alpha,q}^{nc}(s, M) + C_{n\alpha}^c(s, M) \alpha_e \left( \frac{1 + \sqrt{f''}}{2} \right)^2 \quad (1.36)$$

Note that González 2014; Sheng, Galbraith, and Coton 2007 propose the corrective factor to be:

$$C_n^{fs} = C_{n\alpha,q}^{nc}(s, M) + C_{n\alpha}^c(s, M) \alpha_e \left( \frac{1 + 2\sqrt{f''}}{3} \right)^2 \quad (1.37)$$

to account for lower values of  $C_n$  when  $f=0$  that were seen from experimental data.

### 1.2.1 Tangential Force

The along-chord force coefficient analogously becomes:

$$C_c^{fs} = C_c^{pot} \eta_e \left( \sqrt{f''} - 0.2 \right) \quad (1.38)$$

where González 2014's 0.2 factor is used to modify  $\sqrt{f''}$  to account for negative values seen at  $f=0$ .

### 1.2.2 Pitching Moment

Now turning to the pitching moment, the contribution due to unsteady separated flow is on the circulatory component alone. Leishman 2011 suggests using this formulation that modified  $C_m^{pot}$  for the  $C_{m\alpha}^c(s, M)$  component:

$$C_m^{fs} = C_{m0} - C_{n\alpha,q}^c(s, M)(\hat{x}_{cp} - 0.25) + C_{mq}^c(s, M) + C_{m\alpha}^{nc}(s, M) + C_{mq}^{nc}(s, M) \quad (1.39)$$

where  $\hat{x}_{cp}$  is center-of-pressure distance from LE in percent chord and can be approximated by (Leishman 2011):

$$\hat{x}_{cp} = k_0 + k_1(1 - f'') + k_2 \sin(\pi f''^{k_3}) \quad (1.40)$$

where  $k_0=0.25-\hat{x}_{AC}$ , and the  $k_1$ - $k_3$  constants are calculated via best-fits of experimental data. Other expressions could be used to perform the best fit of  $\hat{x}_{cp}$  vs.  $f$  from static  $C_m$  airfoil data.

Minnema 1998 suggests a different approach where an effective lagged AOI is calculated as follows:

$$\begin{aligned} \alpha'_f &= \alpha_f - D_{\alpha f} \\ D_{\alpha f_t} &= D_{\alpha f_{t-1}} \exp\left(-\frac{\Delta s}{0.1T_f}\right) + (\alpha_{f_t} - \alpha_{f_{t-1}}) \exp\left(-\frac{\Delta s}{2*0.1T_f}\right) \end{aligned} \quad (1.41)$$

Note that  $T_f$  is factored by 0.1 following Minnema 1998's validation results. Then the new AOI is used to derive the contribution to the circulatory component of the pitching moment, which is extracted from a look-up table of static coefficients  $C_m$  vs.  $\alpha$ .

$$C_m^{fs} = C_m(\alpha_f') + C_{m_q}^c(s, M) + C_{m_\alpha}^{nc}(s, M) + C_{m_q}^{nc}(s, M) \quad (1.42)$$

The method proposed by González 2014 uses a value of  $f'$  ( $=f_m'$ ) extracted from a static data table where it is assumed that  $C_m - C_{m0} = f_m C_n$  (loosely correlating  $f$  to  $\hat{x}_{cp}$ ). The angle used for interpolation of the lookup table is  $\alpha_f$ . In order to resolve the contribution to  $C_m$  from the unsteady TE separation, this method should then use the full  $C_n^{fs}$  from Eq. (1.37) and the lagged  $=f_m''$  calculated from  $f_m'$  and Eq. (1.34), replacing  $f'$  with  $f_m'$ . However, for the current implementation, there is no further lagging of the  $f_m'$  quantity. This may change in future versions of the UA.

$$C_m^{fs} = C_{m0} + C_n^{fs} f_m' + C_{m_q}^c(s, M) + C_{m_\alpha}^{nc}(s, M) + C_{m_q}^{nc}(s, M) \quad (1.43)$$

In this case, the two treatments (Minnema 1998; González 2014) seem somewhat equivalent, with the exception that González 2014 proposes 21 (7 for each  $f''$  related to  $C_n$ ,  $C_c$ , and  $C_m$ ) different multipliers for  $T_f$  depending on the state of the airfoil aerodynamics (e.g., increasing AOA and above a critical  $C_{n1}$ , increasing AOA and below a critical  $C_{n1}$ ).

### 1.3 Dynamic Stall

During DS there is shear layer roll-up at the LE, vortex formation, and vortex travel over the upper surface of the airfoil to be subsequently shed in the wake. The main condition to be met for the shear layer roll up is:

$$\begin{aligned} C_n' &> C_{n1} & \text{for } \alpha &\geq \alpha_0 \\ C_n' &< C_{n2} & \text{for } \alpha &< \alpha_0 \end{aligned} \quad (1.44)$$

The normal force coefficient contribution from the additional lift associated with the low pressure LE vortex can be written as Leishman 2011:

$$C_{nt}^v = C_{nt-1}^v \exp\left(-\frac{\Delta s}{T_V}\right) + (C_{Vt} - C_{Vt-1}) \exp\left(-\frac{\Delta s}{2T_V}\right) \quad (1.45)$$

$T_V$  is the time constant associated with the vortex lift decay process; it is used in the expression of  $C_n^v$ . It depends on  $Re, M$ , and airfoil class. Note that the  $C_n^v$  contribution is not allowed to have a sign opposite to that of  $C_n^{fs}$ .

$T_V$  gets modified via a multiplier  $\sigma_3$  to account for various stages of the process as discussed later, here suffice to say that:

$$T_V = T_{V0}/\sigma_3 \quad (1.46)$$

$C_V$  represents the contribution to the normal force coefficient due to accumulated vorticity in the LE vortex.  $C_V$  is modeled proportionally to the difference between the attached and separated circulatory contributions to  $C_n$ :

$$C_V = C_{n\alpha,q}^c(s, M) - C_{n\alpha,q}^c(s, M) \left(\frac{1 + \sqrt{f''}}{2}\right)^2 = C_{n\alpha}^c(s, M) \alpha_e \left(1 - \frac{1 + \sqrt{f''}}{2}\right)^2 \quad (1.47)$$

If González 2014 is used, then  $C_V$  writes as:

$$C_V = C_{n\alpha}^c(s, M) \alpha_e \left(1 - \frac{1 + 2\sqrt{f''}}{3}\right)^2 \quad (1.48)$$

If  $\tau_V > T_{VL}$  and if  $\alpha_f$  is not moving away from stall (i.e.,  $(\alpha_f - \alpha_0) * (\alpha_{f_t} - \alpha_{f_{t-1}}) > 0$ ), then the vorticity is no longer allowed to accumulate, in which cases Eq.(1.45) rewrites as:

$$C_{nt}^v = C_{nt-1}^v \exp\left(-\frac{\Delta s}{T_{V0}/\sigma_3}\right) \quad (1.49)$$

with  $\sigma_3 = 2$

where the decay of the normal force (due to vorticity at the LE) is accelerated at twice the original rate and no further accretion of vorticity is allowed. Eq. (1.49) should also be used when conditions in Eq. (1.44) are not met. Note that  $T_{VL}$  represents the time constant associated with the vortex advection process; it represents the non-dimensional time in semi-chords, needed for a vortex to travel from LE to TE; it is used in the expression of  $C_n^v$ . It depends on  $Re$ ,  $M$  (weakly), and airfoil. Value's range = [6; 13].

Finally the total normal force can be written as:

$$C_n = C_n^{fs} + C_n^v = C_{n\alpha}^c(s, M) \alpha_e \left( \frac{1 + \sqrt{f''}}{2} \right)^2 + C_{n\alpha,q}^{nc}(s, M) + C_n^v \quad (1.50)$$

Again, if González 2014 is used, then the correction factor for the separated flow treatment is slightly modified as in Eq.(1.48), i.e.:

$$C_n = C_n^{fs} + C_n^v = C_{n\alpha}^c(s, M) \alpha_e \left( \frac{1 + 2\sqrt{f''}}{3} \right)^2 + C_{n\alpha,q}^{nc}(s, M) + C_n^v \quad (1.50b)$$

Note that multiple vortices can be shed at a given shedding frequency corresponding to:

$$T_{sh} = 2 \frac{1 - f''}{St_{sh}} \quad (1.51)$$

Therefore  $\tau_V$  is reset to 0 if  $\tau_V = T_{VL} + T_{sh}$ .

### 1.3.1 Tangential Force

The along-chord force coefficient gets modified by the presence of the LE vortex as (Pierce 1996):

$$C_c = C_c^{fs} + C_n^v \tan(\alpha_e) \left( 1 - \frac{\tau_V}{T_{VL}} \right) \quad (1.52)$$

Note that in the current release of UA, the  $\tan(\alpha_e) \simeq \alpha_e$  approximation is made. This may be changed after testing in future releases.

González 2014 does not contain the vortex contribution to  $C_c$  based on experimental validation:

$$C_c = C_c^{fs} \quad (1.52b)$$

The original (Leishman and Beddoes 1989) model had  $C_c$  written as:

$$C_c = \begin{cases} \eta_e C_c^{pot} \sqrt{f''} \sin(\alpha_e + \alpha_0) & , \quad C_n' \leq C_{n1} \\ \hat{k}_1 + C_c^{pot} \sqrt{f''} f''^{\hat{k}_2} \sin(\alpha_e + \alpha_0) & , \quad C_n' > C_{n1} \end{cases} \quad (1.53a)$$

$$\text{with : } \hat{k}_2 = 2(C_n' - C_{n1}) + f'' - f \quad (1.53b)$$

where  $\hat{k}_1$  is a constant required to fit the  $C_c$  curve under static conditions for two-dimensional (2D) airfoils.

### 1.3.2 Pitching Moment

Leishman 2011 offers a form for the  $\hat{x}_{cp}^v$ , which is the center-of-pressure distance from the 1/4-chord, in percent chord, during the LE vortex advection process:

$$\begin{aligned} C_m^v &= -\hat{x}_{cp}^v C_n^v \\ \hat{x}_{cp}^v(\tau_V) &= \bar{\bar{x}}_{cp} \left( 1 - \cos \left( \frac{\pi \tau_V}{T_{VL}} \right) \right) \end{aligned} \quad (1.54)$$

where  $\bar{\bar{x}}_{cp}$  is a constant in the expression of  $\hat{x}_{cp}^v$ , usually = 0.2.

Finally, the expression for the total pitching moment can be written as:

$$C_m = C_{m0} - C_{n_{\alpha,q}}^c(s, M)(\hat{x}_{cp} - 0.25) + C_{m_q}^c(s, M) + C_{m_{\alpha}}^{nc}(s, M) + C_{m_q}^{nc}(s, M) + C_m^v \quad (1.55)$$

If Minnema 1998's approach is used then Eq.(1.55) rewrites as:

$$C_m = C_m(\alpha_f) + C_{m_q}^c(s, M) + C_{m_{\alpha}}^{nc}(s, M) + C_{m_q}^{nc}(s, M) + C_m^v \quad (1.56)$$

and if González 2014's treatment is used then the total moment becomes:


$$C_m = C_n f_m'' + C_{m_q}^c(s, M) + C_{m_{\alpha}}^{nc}(s, M) + C_{m_q}^{nc}(s, M) + C_m^v \quad (1.57)$$



## 2 Inputs, Outputs, Parameters, States and Implementation of the UA

The UA implementation loosely follows the FAST modularization framework. This includes key interface routines for: module initialization (UA\_Init), updating of the module's states (UA\_UpdateStates), and generating module outputs (UA\_CalcOutput). Since this module does not communicate directly with the FAST glue code, we have taken some liberties with regards to the arguments to this interface routines. Additionally, the typical framework data structures for inputs (u) and outputs (y) have been modified to allow for a more intuitive integration into either a BEMT algorithm or the GDW algorithm. In the following sections, first the inputs/states/parameters are listed, and then each interface routine will be discussed and any deviations from the FAST framework will be highlighted.

In order to provide the context for the UA sub-module, flow-diagram from AeroDyn to UA is shown in Fig. 4:



**Figure 4. Block diagram showing the order of the calls to the subroutines (1st and 2nd in the graph) and the overall organization from the parent module AeroDyn to UA.**

### 2.1 Init\_Inputs

Beside the standard variables common to all modules (*OutFmt*, *OutSFmt*, *NumOuts*, *OutList*), the Init\_Inputs to the UA are:

- $\Delta t$  -time step
- $a_s$  -speed of sound
- *DSMod* - switch to select handling of options and possible methods in dynamic stall
- *AFIParams* -airfoil static tables of  $C_l$   $C_d$   $C_m$  and DS parameters
- *AFidx* -index pointing to the correct airfoil file/table/database

- $c$  - airfoil chord at each blade station for each blade.

## 2.2 Inputs $u$

The Inputs to the UA are:

- $\alpha$
- $U$
- $Re$

Note: these are for a given node (within a given blade) and chord.

## 2.3 Outputs $y$

The Outputs from the UA are:

- $C_n$
- $C_c$
- $C_m$
- $C_l$
- $C_d$

Note: these are for a given node (within a given blade) and chord.

## 2.4 States $x_d$

The States for the UA are:

*Discrete States:*

- $\alpha_{-1}$  -previous time-step value of  $\alpha$
- $\alpha_{-2}$  -previous time-step value of  $\alpha$
- $q_{-1}$  -previous time-step value of  $q$
- $q_{-2}$  -two time-steps ago value of  $q$
- $X_{1,-1}$  -previous time-step value of  $X_1$
- $X_{2,-1}$  -previous time-step value of  $X_2$
- $X_{3,-1}$  -previous time-step value of  $X_3$
- $X_{4,-1}$  -previous time-step value of  $X_4$
- $K'_{\alpha,-1}$  -previous time-step value of  $K'_{\alpha}$
- $K'_{q,-1}$  -previous time-step value of  $K'_q$
- $K''_{q,-1}$  -previous time-step value of  $K''_q$
- $K'''_{q,-1}$  -previous time-step value of  $K'''_q$
- $D_{p,-1}$  -previous time-step value of  $D_p$

- $D_{f,-1}$  -previous time-step value of  $D_f$
- $C_n^{pot,-1}$  -previous time-step value of  $C_n^{pot}$
- $T_f$  -constant dependent on Mach,  $Re$ , and airfoil shape; it is used in the expression of  $D_f$  and  $f''$
- $f'_{-1}$  -previous time-step value of  $f'$
- $f''_{-1}$  -previous time-step value of  $f''$
- $\tau_V$  -time variable, tracking the travel of the LE vortex over the airfoil suction surface. It is made dimensionless via the semi-chord:  $\tau_V = t * 2U / c$ . If less than  $2T_{VL}$ , it renders  $VRTX=True$ ; if less than  $T_{VL}$  then the vortex is still on the airfoil.
- $C_{n,-1}^v$  -previous time-step value of  $C_n^v$
- $C_{V,-1}$  -previous time-step value of  $C_V$
- $D_{\alpha f,-1}$  -previous time-step value of  $D_{\alpha f}$

The FAST 8 framework does not allow logicals or discontinuous variables within states. For this reason, the following ones are declared as Other States.

*Other States:*

- $\sigma_1$  -generic multiplier for  $T_f$
- $\sigma_3$  -generic multiplier for  $T_V$
- $TESF$  -trailing edge separation flag
- $LESF$  -leading edge separation flag
- $VRTX$  -vortex advection flag
- $FirstPass$  -flag indicating first time step

Note that, in contrast to Inputs and Outputs, the States must be tracked by the UA module, therefore they are two-dimensional array (per blade, per node).

## 2.5 Parameters p

The Parameters for the UA are :

- $\Delta t$
- $c$
- $DSMod$
- $a_s$
- $AFIParams$
- $AFidx$

The airfoil  $AFIParams$  structure contains airfoil specific quantities, i.e. parameters and constants for the UA:

- $\alpha_0, \alpha_1, \alpha_2, C_{n\alpha}, C_{n1}, C_{n2}, \eta_e, C_{d0}, C_{m0}, \bar{x}_{cp}, St_{sh}$
- $A_1, b_1, A_2, b_2, A_5, b_5$
- $S_1, S_2, S_3, S_4$

- $T_p$ , fairly independent of airfoil type
- $T_{f0}, T_{V0}, T_{VL}$
- $k_0, k_1, k_2, k_3$
- $\hat{k}_1$

## 2.6 UA Implementation

### 2.6.1 UA\_Init Routine

This routine allocates the module's data structures, initializes the module's states, and sets the non-time-varying parameters (copies them from the initialization input data section).

### 2.6.2 UA\_UpdateStates Routine

The typical list of arguments to UA\_UpdateStates gets augmented to pass indices to the blade and blade-node of interest. The indices point into the array of structures  $AFI_{Params}$  and into the array of chords.

The model is of the parsimonious, open loop, Kelvin-chain kind. Outputs of one subsystems go into inputs of the next subsystems. There are no differential equations to solve. There is no solver per se, for this reason states are discrete states only (and other states see Section 2.4).

#### 2.6.2.1 DSMod Logical Flags

The options implemented in the code are as follows:

IF Gonzalez THEN

- Replace Eq. (1.55) with Eq. (1.57)
- Replace Eq. (1.52) with Eq. (1.52b)
- Replace Eq. (1.50) with Eq. (1.50b)
- Replace Eq. (1.47) with Eq. (1.48)
- Replace Eq. (1.39) with Eq. (1.43)
- Replace Eq. (1.36) with Eq. (1.37)
- Add Eq. (1.17) to  $C_{n\alpha,q}^c(s, M)$  Eq. (1.14)

IF Minemma THEN

- Replace Eq. (1.55) with Eq. (1.42)
- Replace Eq. (1.39) with Eq. (1.41)-(1.42)
- Replace Eq. (1.27) with Eq. (1.28)

IF flookup THEN

- Replace Eq. (1.31) with expressions for  $f_n' f_n'$
- IF Gonzalez THEN calculate  $f_m''$

IF CCLBMswitch THEN

- Replace Eq. (1.52) with Eq. (1.53)

#### 2.6.2.2 Update Discrete States

For a given set of inputs (u), and at the current step in time (t), the Kelvin chain is performed through the following equations:

- Eq. 1.12c
- Eq. 1.5b
- Eq. (1.8)
- Eq. (1.12a)
- Eq. (1.12b)
- Eq. (1.11)
- Eq. (1.35)
- Eq. (1.46)
- Eq. (1.19)
- Eq. (1.20)
- Eq. (1.18)
- Eq. (1.16)
- Eq. (1.15)
- Eq. (1.14)
- Eq. (1.23)
- Eq. (1.22)
- Eq. (1.21)
- Eq. (1.26)
- Eq. (1.29)
- Eq. (1.33)
- Eq. (1.32)
- Eq. (1.31)
- Eq. (1.34)
- Eq. (1.36) (or Eq. (1.37))
- Eq. (1.47) (or Eq. (1.48))
- Eq. (1.45) (or Eq. (1.49))

### 2.6.2.3 Update Other States

- IF  $C'_n > C_{n1}$  ( $C'_n < C_{n2}$  for  $\alpha < \alpha_0$ ) THEN  
 $LESF=TRUE$ : this means LE separation can occur.  
ELSE  $LESF=False$ : this means reattachment can occur.
- IF  $f''_t < f''_{t-1}$  THEN  
 $TESF=True$ : this means TE separation is in progress.  
ELSE  $TESF=False$ : this means TE reattachment is in progress.

- IF  $0 < \tau_V \leq 2T_{VL}$  THEN  
 $VRTX=True$ : this means vortex advection is in progress.  
 $ELSE VRTX=False$ : this means vortex is in wake.
- IF  $\tau_V \geq 1 + \frac{T_{sh}}{T_{VL}}$  and  $LESF=True$  THEN  
 $\tau_V$  is reset to 0.

### 2.6.2.3.1 $T_f$ modifications

The following conditional statements operate on a multiplier  $\sigma_1$  that affects  $T_f$ , i.e., the actual  $T_f$  is given by Eq.(1.35).

$$T_f = T_{f0}/\sigma_1 \quad (1.35 \text{ revisited})$$

where  $T_{f0}$  is the initial value of  $T_f$ .

$\sigma_1 = 1$  (initialization default value)

$\Delta\alpha_0 = \alpha - \alpha_0$

IF  $TESF=True$  THEN: (separation)

IF  $K_\alpha \Delta\alpha_0 < 0$  THEN  $\sigma_1 = 2$  (accelerate separation point movement)

ELSEIF  $LESF=False$  THEN  $\sigma_1 = 1$  (default value, LE separation can occur)

ELSEIF  $f''_{n-1} \leq 0.7$  THEN  $\sigma_1 = 2$  (accelerate separation point movement if separation is occurring)

ELSE  $\sigma_1 = 1.75$  (accelerate separation point movement)

ELSE: (reattachment, this means  $TESF=False$ )

IF  $LESF = False$  THEN  $\sigma_1 = 0.5$  (default: slow down reattachment)

IF  $VRTX=True$  AND  $0 \leq \tau_V \leq T_{VL}$  THEN

$\sigma_1 = 0.25$  - No flow reattachment if vortex shedding in progress

IF  $K_\alpha \Delta\alpha_0 > 0$  THEN  $\sigma_1 = 0.75$

Note the last three conditional statements are separate IFs.

While this logic was tested and proved to be effective, the current version of UA uses a simpler version.

### 2.6.2.3.2 $T_V$ modifications

For  $T_V$ , an analogous set of conditions is used to set the proper value of the time constant depending on subsystem stages:

$\sigma_3=1$  (initialization default value)

IF  $T_{VL} \leq \tau_V \leq 2T_{VL}$  THEN

$\sigma_3=3$  (post-shedding)

IF  $TESF=False$  THEN

$\sigma_3=4$  (accelerate vortex lift decay)

IF  $VRTX=True$  AND  $0 \leq \tau_V \leq T_{VL}$  THEN:

IF  $K_\alpha \Delta\alpha_0 < 0$  THEN  $\sigma_3=2$  (accelerate vortex lift decay)

ELSE  $\sigma_3=1$  (default)

ELSEIF  $K_\alpha \Delta\alpha_0 < 0$  THEN  $\sigma_3=4$  (vortex lift must decay fast)

IF  $TESF=False$  AND  $K_q \Delta\alpha_0 < 0$  THEN  $\sigma_3=1$  (default)

### 2.6.2.3.3 Update 'previous time step' states

After the states are updated to the next time step (t+1) values, the current values at time step t, are stored into the (t-1) states.

### 2.6.3 *UA\_CalcOutput*

This routine determines the outputs,  $C_n$ ,  $C_c$ ,  $C_t$  (and the transformed versions,  $C_l$  and  $C_d$ ) given the inputs of  $U$ ,  $\alpha$ , and  $Re$  (currently unused), for a given blade element. Since the routine only generates outputs for a specific blade element, the FAST framework arguments are augmented to include indices to the blade and blade-node of interest. The routine uses the same Kelvin chain as in *UA\_UpdateStates*, however, the state variables themselves are not updated during these calculations.

For the first time step, outputs are determined by static lookup tables.

The equations implemented in this routine are the same as in Section 2.6.2.2, plus the following ones:

- Eq. (1.50)
- Eq. (1.52)
- Eq. (1.2)
- Eq. (1.55)

## Bibliography

- Damiani, R. 2011. *Algorithmic Outline of Unsteady Aerodynamics (AeroDyn) Modules*. Final Report WE-201103. NREL Subcontract No. AFT-1-11326-01 under Prime Contract No. DE-AC36-08GO28308. Available upon request from NREL. Arvada, CO: RRD Engineering, LLC.
- González, A. 2014. *DYSTOOL Stability Aerodynamic Tool for 2D Airfoil Based on the Beddoes-Leishman Model*. Informe IN-523/069-en. Wind Energy Department – Fundación CENER-CIEMAT. Avenida Ciudad de la Innovación, 7 - 31621-Sarriguren (Navarra) Spain: CENER.
- Gupta, S., and J. G. Leishman. 2006. “Dynamic Stall Modeling of the S809 Airfoil and Comparison with Experiments”. *Wind Energy* 9 (6): 521–547.
- Johansen, J. 1999. *Unsteady Airfoil Flows with Application to Aeroelastic Stability*. Report Risø-R-1116(EN). 98 pages. Roskilde, Denmark: Risø National Laboratory.
- Jonkman, J. 2013. “New Modularization Framework for the FAST Wind Turbine CAE Tool”. In *Proceedings of the 51st AIAA Aerospace Sciences Meeting*. Dallas, TX: AIAA.
- Leishman, J. G. 2006. *Principles of Helicopter Aerodynamics*. 2nd. 826 p. New York, NY: Cambridge University Press.
- . 2011. *Final Report: Assessment of 'AeroDyn' Theory Basis Including Unsteady Aerodynamics Modules*. NREL Subcontractor Report LFC-1-11303-01. Prime Contract No. DE-AC36-08GO28308.
- Leishman, J. G., and T. S. Beddoes. 1986. “A General Model for Airfoil Unsteady Behavior and Dynamic Stall Using the Indicial Method”. In *Proceedings of the 42nd Annual Forum of the American Helicopter Society*, 243–266. Washington, D.C.: American Helicopter Society.
- . 1989. “A Semi-Empirical Model for Dynamic Stall”. *J. of the American Helicopter Society* 34 (3): 3–17.
- Lomax, H., et al. 1952. *Two and Three Dimensional Unsteady Lift Problems in High Speed Flight*. Technical Report 1077. NACA.
- Minnema, J. E. 1998. *Pitching Moment Predictions on Wind Turbine Blades Using the Beddoes-Leishman Model for Unsteady Aerodynamics and Dynamic Stall*. MA thesis, The University of Utah.
- Pierce, K. G. 1996. *Wind Turbine Load Prediction Using the Beddoes-Leishman Model for Unsteady Aerodynamics and Dynamic Stall*. MA thesis, The University of Utah.
- Pierce, K. G., and A. C. Hansen. 1995. “Prediction of Wind Turbine Rotor Loads Using the Beddoes-Leishman Model for Dynamic Stall”. *J. of Solar Energy Engineering* 117:200–204.
- Sheng, W., R. A. Galbraith, and F. N. Cotton. 2007. “A Modified Dynamic Stall Model for Low Mach Numbers”. In *45th AIAA Aerospace Sciences Meeting and Exhibit*.